Crisis Response Service Standards

for Mental Health Services and Supports

May, 2005



Table of Contents

2 2 2	Introduction Crisis Response Services
3	Need for Crisis Response Service Standards Broad Policy Context
• •	broad Foncy Context
4	Provincial Policy Context
4	Making It Happen
4	Mental Health Accountability Framework
5	Process for Developing Standards
5	Interjurisdictional Review
5	Stakeholder Working Group
6	Validation of Standards
6	Crisis Response Service Standards
6	Features of Crisis Response Services - A Vital Component of
	the Service Continuum
6	Crisis Response Service Functions
7	Relationship Between Service Standards, Functions, Domains, Indicators and Performance Measures
8	Crisis Response Service Standards
12	Next Steps
13	Appendix A: Performance Domains
14	Appendix B: Mental Health Accountability Framework -
	Performance Domains and Indicators
16	Appendix C: List of Stakeholder Working Group Participants
17	Appendix D: Functions of Crisis Response Services
18	Appendix E: Crisis Response Service Standards, Domains and Indicators
30	Appendix F: Literature Review – Summary of Key Research Findings on Crisis Response Services
39	Appendix G: Bibliography

Validation of Standards

The set of standards drafted by the stakeholder working group was validated through a review of academic literature and a scan of crisis response service standards in provinces across Canada. A "levels of evidence" typology was based on a typology developed by the Nova Scotia Department of Health (Nova Scotia Department of Health, 2003) (See Appendix F for a description of the levels of evidence typology) to compare the proposed standards with available evidence. Relevant literature was reviewed against the standards. Each standard was rated based on the levels of evidence typology and evidence supporting the standard was identified.

Crisis Response Service Standards

Features of Crisis Response Services – A Vital Component of the Service Continuum

Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports describes the features of comprehensive crisis response services.

A crisis is defined as the onset of an emotional disturbance or situation distress (which may be cumulative) involving a sudden breakdown of an individual's ability to cope.

Crisis response services are part of an integrated system of mental health services and should provide timely access to a wide range of crisis options on a 24 hour basis. Active treatment and support is offered in a variety of environments as soon as possible after an individual is identified as in acute distress. Services should provide immediate relief of symptoms and rapid stabilization so the condition does not worsen. Crisis response services also offer the opportunity to develop longer-term treatment and rehabilitation plans and have the potential to mobilize community resources and avert the need for short and/or long-term hospitalization.

Examples of crisis response services include telephone crisis response, walk-in services, mobile crisis outreach, crisis residential services and psychiatric emergency/medical crisis services.

Crisis response services are available to all people with symptoms of mental illness and take into account the age, gender, race, language, etc., of the person. Priority is given to individuals with serious mental illness and those who may be at risk of causing harm to themselves or others.

Crisis Response Service Functions

A range of services and supports are required by consumers and their families to assist in crisis prevention and on-going support. *Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports* sets out the following specific functions of a crisis service. (See Appendix D for a definition of the functions.)


- Assessment and Planning
- Crisis Support/Counseling
- Medical Intervention
- Environmental Interventions and Crisis Stabilization
- Review/Follow-up/Referral
- Monitoring and Evaluation
- Information, Liaison, Advocacy and Consultation/Collaboration

Standards were developed for each of the crisis response service functions and reflect the general features of crisis response services.

Relationship Between Service Standards, Functions, Domains, Indicators and Performance Measures

Compliance with standards will ensure services are comprehensive, coordinated and based on consumer need and best practices. The crisis response service standards reflect the key features and functions of crisis response services laid out in *Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports*, and the performance domains and indicators identified in the *Mental Health Accountability Framework*.

Figure 1: Relationship Between Service Standards, Functions, Domains, Indicators and Performance Measures



Each crisis response service standard relates to one or more of the domains and indicators.

The next step will be the development of performance measures based on the applicable domains and indicators. For this reason, every standard is associated with domain(s) and indicators and has been worded so it can be translated into measurable statements. (Refer to Appendix E for a table presenting the standards and the related domains and indicators.) Each indicator may encompass a number of measurements relevant to the service provided. For example:

Indicator: Accessibility – Wait times for needed services.

Standard: Upon identification of a crisis, the first contact with the consumer by the crisis response service must be established within 90 minutes.

Measure:

Number of consumers where first contact was established in:

Under 15 minutes

Between 16 and 30 minutes

Between 31 and 45 minutes

Between 46 and 60 minutes

Between 61 and 75 minutes

Between 76 and 90 minutes

Over 90 minutes

• Indicator: Acceptability - Consumer/family satisfaction with service provided.

Standard: A consumer satisfaction survey should be developed and implemented on an annual basis, with a target of 80% satisfaction, to the best extent possible, given the mandate of the CRS.

Measure:

Percentage of consumers who reported in the consumer satisfaction survey that they were satisfied with the service received.

Less than 50%

51-60%

61-70%

71-80%

81-90%

91-100%

Once performance measures are developed for each indicator related to a crisis response standard, data will be collected and used to evaluate services and further refine the standards as required. Standards and services will continually be re-evaluated based on the measures and collected data to ensure the standards reflect best practices.

Crisis Response Service Standards

The standards presented in this document have been developed based on the service functions in *Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports*, and the domains and indicators defined in the *Mental Health Accountability Framework*. They also reflect consultation with the stakeholder working group, developments in other jurisdictions and research evidence.

The following chart presents each standard, its related service function and domain(s) and identifies the level of evidence (i.e., research evidence, expert opinion) supporting the standard. (See Appendix F for a description of the levels of evidence typology.)

Table 1: Crisis Response Service Standards

				evel vide		
Function	Standard	Domain	1	2	3	4
Assessment and Planning	 Upon identification of a crisis, the first contact with the consumer by the crisis response service (CRS) must be established within 90 minutes. 	Accessibility				✓
	 A crisis requiring in-person contact will be responded to as soon as possible. Response time should be within 24 hours, with consideration for travel time, weather, etc. 	Accessibility				1
	 Crisis support telephone lines must be configured to include a queuing system that lasts no longer than 15 minutes. During the wait time, voice instructions for alternative crisis management options must be provided (e.g., 911, local emergency number, another crisis line). 	Accessibility				✓
	 Protocols must be in place with related service providers (e.g., case management services, psychiatrists, hospitals, primary care teams, etc.) in order to ensure access to necessary medical, psychiatric and psychological/social assessments and existing crisis management protocols. If access to other health records will assist in planning with the consumer, documented consent is required. 	Accessibility		1		1
	Services must be provided in the consumer's place of choice wherever possible.	Accessibility			,	7

Levels of Evidence

Level 1 involved direct evidence of effectiveness (i.e., specific standard evaluated as independent variable in a study and shown to produce positive outcomes).

Level 2 considered indirect evidence of effectiveness (i.e., specific standard is one of the characteristics or ingredients of a program which has been shown to be effective).

Level 3 involved studies based on expert opinion/consensus of effectiveness or correlational evidence of standards being associated with positive consumer outcomes.

Level 4 involved expert opinion (i.e., defined by the working group or other Canadian jurisdictions in absence of empirical support in research literature).

Table 1: continued

Assessment and Planning (continued)	 The CRS will ensure that staff have training and core competencies in a variety of areas, to the best extent possible, (e.g., risk assessment; suicide assessment; crisis intervention; safety standards; conflict resolution; anti-racism training; psychiatric symptomatology and psychiatric medications). 	Competence	
	 The CRS must ensure that workers are provided with information regarding relevant legislative reporting requirements. 	Competence	\
Crisis Support/ Counseling	 Services are consumer directed and will be provided in the least intrusive manner possible. 	Accessibility, Acceptability and Appropriateness	
	 Written protocols for consumer and service provider safety must be established. 	Safety	1
	 Written protocols must be developed to ensure that timely consultation is available with various service providers (e.g., physicians, guidance counselors, CAS workers) to assist in identifying and intervening in actual emergencies; documented consent is required. 	Accessibility	
	 Short-term crisis support/counseling will be available to provide risk assessment, de-escalation and safety planning, to the best extent possible, given the mandate of the CRS. 	Accessibility	1
Medical Intervention	 The CRS must have written protocols for initiating and accessing medical interventions in a timely manner. 	Accessibility	1
	 The CRS will have procedures/ guidelines for responding to emergency and non-emergency medical situations. 	Appropriateness	1
Environmental Interventions and Crisis Stabilization	All CRS will have access to other services 24 hours a day, seven days a week.	Accessibility	1
подысывает	All CRS will have access to current community contact information.	Competence	1

Table 1: continued

Environmental Interventions and Crisis Stabilization (continued)	 Protocols are in place for providing referral and support (i.e., income support, dealing with employers, access to long/short-term housing, addressing family issues) based on consumer-articulated needs. 	Appropriateness and Accessibility	1
Review/ Follow-up/ Referral	The consumer must have the opportunity to review, discuss and comment on the service and its appropriateness.	Appropriateness and Acceptability	1
	 A review process must be established that includes an assessment of consumer outcomes including status and stability (e.g., Did the consumer perceive return of control? Was the crisis stabilized?) 	Effectiveness	1
	Written protocols must be established for providing referral and transition to post-crisis services. Referrals to post-crisis services must be based on consumer-articulated needs.	Appropriateness	√
	CRS must ensure that staff are provided with information about other relevant community resources.	Competence	✓
	 A written follow-up plan must be developed upon completion of service, which will include criteria for follow-up, re-entry and linkage with other services. 	Effectiveness and Acceptability	✓
Monitoring and Evaluation	 Consumer satisfaction (including consumers, families and outside agencies) must be monitored continuously, and the results used to make service improvements. 	Acceptability	√
	 A consumer satisfaction survey will be developed and implemented on an annual basis, with a target of 80% satisfaction, to the best extent possible, given the mandate of the CRS. 	Acceptability	√
	 All organizations and agencies must evaluate some aspect of their programs annually using best practices and published standards. 	Effectiveness and Appropriateness	1

Table 1: continued

Information, Liaison, Advocacy, and Consultation/ Collaboration	 Service providers must identify gaps in service and develop means for collaborating with other relevant community resources in order to meet unmet needs. 	Efficiency and Continuity	1
	 A plan must be developed and published that identifies community linkages. 	Effectiveness	1
	 The crisis worker must be knowledgeable about services that are accessible and relevant to consumers' interests in order to provide up-to-date information. 	Competence and Appropriateness	1
	 CRS must ensure that staff is provided with information about other relevant community resources. 	Effectiveness	V

Next Steps

This document describes crisis response service standards and represents the next step in the development of accountable crisis response services in Ontario. The ministry will be further developing the components of the *Mental Health Accountability Framework* in order to define and implement an accountability relationship and process with the mental health system.

The principles, goals and essential components of a reformed mental health system have been defined. Performance domains, indicators and standards have been developed, based on system principles and goals and program components. This represents one step in the provision of clear program direction for crisis response services to the field. While all service providers may not be able to meet all standards immediately, significant funding is being invested in the enhancement of crisis response services to ensure that services will be able to meet standards.

The ministry will also develop data collection requirements and outcome-based performance measures to monitor and report on the provision of crisis response services. As these performance measures and data collection requirements are implemented, the available data will be used to measure crisis response services and supports against the identified standards.

With these components in place, the ministry will be able to implement an accountability process to monitor how crisis response services are being provided and answer important questions.

Are services being delivered across the province in a manner that is consistent with ministry policy and with evidence-based best practices?

Most importantly, are consumers satisfied with the service they are receiving and is the service helping them to achieve their personal goals?

This will inform the continual improvement and evaluation of the system of crisis response services within Ontario's mental health system.

Appendix A: Performance Domains

Domain	Definition	
Acceptability	Services provided meet expectations of service users, community, providers and government.	
Accessibility Ability of people to obtain services at the right place and right time on needs.		
Appropriateness	Services provided are relevant to service user needs and based on established standards.	
Competence	Knowledge, skills and actions of individuals providing services are appropriate to service provided.	
Continuity The system is sustainable, comprehensive, and has the capacity seamless and coordinated services across programs, practitioner organizations and levels of service in accordance with individual		
Effectiveness	Services, intervention or actions achieve desired results.	
Efficiency Organizations/programs achieve desired results with the most cost-effective use of resources.		
Safety	Organizations/programs avoid or minimize potential risks or harms to consumers, families, mental health staff and the community associated with the intervention/lack of intervention or the environment.	

Ontario Ministry of Health and Long-Term Care (2003), Mental Health Accountability Framework, p. 18

Appendix B: Mental Health Accountability Framework -Performance Domains and Indicators'

1 ACTION OF THE PARTY OF THE PA	DO	MAIN	
Acceptability	Accessibility	Appropriateness	Competence
Consumer/family satisfaction with service received	Service reach to persons with serious mental illness (SMI)	Existence of best practice core programs	Resources available to train staff to meet required competencies for role
Consumer/family involvement in treatment decisions	Service reach to the homeless	Fidelity: adherence to best practices	Resources available for on-the-job developmen and continuous learnin
Formal complaints mechanisms in place	Access to psychiatrists and other mental health professionals	Best practices services/ supports provided to persons with SMI	Meets provincial certification/ professional standards (where applicable)
Patient bill of rights	Identify human resource gaps	Treatment protocols for co-morbidity	
Consumer/family involvement in service delivery and planning	Access to primary care	Hospital readmission rate*	
Cultural sensitivity	Wait times for needed services	Involuntary committal rate*	
Consumer/family choice of services	Availability of after hours care	Length of stay in acute care*	
	Availability of transportation	Time in community programs	
	Denial of service	Use of seclusion/ restraints	
	Early intervention	Level of service and setting appropriate to needs of individual	
	Consumer/family perception of accessibility	Needs-based funding and spending	
	Access to continuum of mental health service	Consumer/family perception of appropriateness	
	Criminal justice system involvement	Availability of community services	
		Criminal justice system involvement	
		Community/institutional balance	

D	A	RAT		IN	7
v	v	171	А	ш	ч

DUMAIN			
Continuity	Effectiveness	Efficiency	Safety
Continuity mechanisms	Community tenure	Mental health spending per capita	Complications associated with electro-convulsive therapy (ECT)
Emergency room visits*	Mortality	Proportion of staff funding spent on administration and support	Medication errors
Community follow-up after hospitalization	Criminal justice system involvement	Needs-based allocation strategy	Medication side effects
Documented discharge plans	Clinical status	Community/institutional balance	Critical incidents
Cases lost to follow-up	Functional status	Resource intensity planning tool	Suicides
Clear, visible and available points of accountability	Involvement in meaningful daytime activity	Unit costs and cost per consumer	Homicides
	Housing status	Budget and tools for evaluation and performance monitoring	Involuntary committal rate
	Quality of life		Risk management practised
	Physical health status		Identify research/ practices to reduce adverse events and errors
			· ·
NOTE: Indicators marked with an aster	risk are often used as measures. They are		

NOTE: Indicators marked with an asterisk are often used as measures. They are included here as indicators to reflect that they may signal system function or problems.

Appendix C: List of Stakeholder Working Group Participants

Anne Bowlby

Manager, Mental Health and Addiction Branch, Community Health Division, MOHLTC

Amy Churchman

Program Manager – Crisis, CMHA Lambton County, Sarnia

Karen D'Alessio

Coordinator, Mobile Crisis Team, St. Elizabeth Health Care, Mississauga

Henry deSouza

Program Manager, East Region, Community Health Division, MOHLTC

Lana Frado

Executive Director, Sound Times Support Services, Toronto

Catherine Ford

Senior Program Analyst, Mental Health and Addiction Branch, Community Health Division, MOHLTC

Vicente Gannam

Senior Addiction Program Analyst, Mental Health and Addiction Branch, Community Health Division, MOHLTC

Sheila Irvine

Administrative Director, Elm Grove, Brockville Site, Royal Ottawa Hospital, Brockville

Steve Keeble

310-Cope (Crisis Program) York Support Services Network, Newmarket

Paul Links

St. Michael's Hospital Arthur Sommer Rotenberg Chair in Suicide Studies, Toronto

Susan Marshall

Can-Help, Fort Frances

Alan Mathany

Director,
Crisis and Community
Support Teams
Frontenac Community
Mental Health Services,
Kingston

Beth McCarthy-Kent

Mental Health Consultant, North Region, Community Health Division, MOHLTC

Pierina Minna

Mental Health Consultant, Central West Region, Community Health Division, MOHLTC

Christine Nichols

Program Manager, Mental Health Crisis Line, Royal Victoria Hospital, Barrie

Julie Pawlick

Program Manager, COAST – Crisis Outreach and Support Team, Hamilton

Paul Quinn

Executive Director, Gerstein Centre, Toronto

Geoff Reekie

Area Program Manager – Crisis, Muskoka-Parry Sound Community Mental Health Service, Bracebridge

Deb Sherman

Mental Health Rights Coalition, Hamilton

Ann Smithson

Team Leader, Community Crisis Care, Niagara Health System, St. Catharines Site

Kyla Storry

Coordinator, Kenora Rainy River District Crisis Response Services, Dryden

Joanne Walsh

Manager, Crisis Services, Inner City Health Program, St. Michael's Hospital, Toronto

Brent Woodford

Executive Director, Adult Mental Health Services of Haldimand-Norfolk, Simcoe

Bernadette Wren

Manager, Community Mental Health Services – Renfrew County, Pembroke

Staff Support

Ruth Stoddart/ Nancy Douglas/Aimee Watson, Mental Health and Rehabilitation Policy Unit, MOHLTC

Appendix D: Functions of Crisis Response Services

Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports (1999) defined the functions of a crisis response service.

- Assessment and Planning includes gathering pertinent information from the consumer and
 other key supports to develop an understanding of recent events, and psychosocial and biological
 factors related to the presenting crisis. This function also includes the development of an intervention
 plan which takes into account the consumer's immediate needs, strengths, weaknesses and social
 support system.
- Crisis Support/Counseling provides the individual and family with emotional support, practical
 assistance and access to a range of appropriate resources available to resolve the immediate crisis.
- Medical Intervention is an integral part of the crisis response system. It is important to develop links between medical and non-medical service providers to ensure access to resources to resolve the crisis. Medical interventions may be provided by nurses, physicians and pharmacists.
- Environmental Interventions and Crisis Stabilization involves access to required services to stabilize the crisis and includes direct action within the individual's community to provide supports such as arranging for money/income support, dealing with employers, planning for long/short-term housing/accommodation issues and addressing family issues.
- Review/Follow-up/Referral provides appropriate referral to ongoing services and supports that have been mutually defined by the consumer and service provider once the crisis has dissipated.
- Monitoring and Evaluation evaluates the achievement of goals (from the perspective of both the consumer and service provider) and consumer satisfaction.
- Information, Liaison, Advocacy and Consultation/Collaboration provides information to the consumer, family/key supports and service providers regarding types of services and supports available. Works to establish partnerships among service providers to create an integrated service network, and advocates and consults on behalf of consumers and families/key supports within the service network.

Appendix E: Crisis Response Service Standards, Domains and Indicators'

Function	Standard
Assessment and Planning	 Upon identification of a crisis, the first contact with the consumer by the crisis response service (CRS) must be established within 90 minutes.
	A crisis requiring in-person contact will be responded to as soon as possible. Response time should be within 24 hours, with consideration for travel time, weather, etc.
	 Crisis support telephone lines must be configured to include a queuing system that lasts no longer than 15 minutes. During the wait time, voice instructions for alternative crisis management options must be provided (e.g., 911, local emergency number, another crisis line).
	 Protocols must be in place with related service providers (e.g., case management services, psychiatrists, hospitals, primary care teams, etc.) in order to ensure access to necessary medical, psychiatric and psychological/social assessments and existing crisis management protocols. If access to other health records will assist in planning with the consumer, documented consent is required.
	Services must be provided in the consumer's place of choice wherever possible.
	• The CRS will ensure that staff have training and core competencies in a variety of areas, to the best extent possible, (e.g., risk assessment; suicide assessment; crisis intervention; safety standards; conflict resolution; anti-racism training; psychiatric symptomatology and psychiatric medications).
	The CRS must ensure that workers are provided with information regarding relevant legislative reporting requirements.

Based on the domains and indicators defined in Mental Health Accountability Framework (Ontario Ministry of Health and Long-Term Care, 2003).

	Domain	Indicators
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to the homeless Wait times for needed services Availability of after hours care Availability of transportation Denial of service
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to the homeless Wait times for needed services Availability of after hours care Availability of transportation Denial of service
	Accessibility	 Service reach to persons with serious mental illness (SMI) Wait times for needed services Availability of after hours care Denial of service
	Accessibility	 Service reach to persons with serious mental illness (SMI) Access to psychiatrists and other mental health professionals Wait times for needed services Access to primary care Availability of after hours care Access to continuum of mental health service Criminal justice system involvement Denial of service
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to the homeless Access to psychiatrists and other mental health professionals Availability of transportation Consumer/family perception of accessibility Access to continuum of mental health service Criminal justice system involvement
	Competence	 Resources available to train staff to meet required competencies for role Resources available for on-the-job development and continuous learning Meets provincial certification/professional standards (where applicable)
	Competence	 Resources available to train staff to meet required competencies for role Resources available for on-the-job development and continuous learning

Function	Standard
Crisis Support/ Counseling	Services are consumer-directed and will be provided in the least intrusive manner possible.
	Written protocols for consumer and service provider safety must be established.
	Written protocols must be developed to ensure that timely consultation is available with various service providers (e.g., physicians, guidance counselors, CAS workers) to assist in identifying and intervening in actual emergencies; documented consent is required.
	Short-term crisis support/counseling will be available to provide risk assessment, de-escalation and safety planning, to the best extent possible, given the mandate of the CRS.

	Domain.	Indicators	
	Accessibility	 Consumer/family perception of accessibility Access to continuum of mental health service 	
	Acceptability	 Consumer/family involvement in treatment decisions Consumer family involvement in service delivery and planning Consumer/family choice of services Cultural sensitivity 	
	Appropriateness	 Best practices services/supports provided to persons with SMI Level of service and setting appropriate to needs of individual Consumer/family perception of appropriateness Availability of community services Community/institutional balance 	
	Safety	 Critical incidents Suicides Homicides Involuntary committal rate Risk management practised Identify research/practices to reduce adverse events and errors Medication side effects 	
	Accessibility	 Access to psychiatrists and other mental health professionals Access to primary care Wait times for needed services Availability of after hours care Availability of transportation Access to continuum of mental health service Criminal justice system involvement Denial of service 	
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to homeless Access to psychiatrists and other mental health professionals Access to primary care Wait times for needed services Availability of after hours care Availability of transportation Consumer/family perception of accessibility Access to continuum of mental health service Criminal justice system involvement Denial of service 	

Function	Standard
Medical Intervention The CRS must have written protocols for initiating and accessing medical interventions in a timely manner.	
	The CRS will have procedures/guidelines for responding to emergency and non-emergency medical situations.
Environmental Interventions and Crisis Stabilization	All CRS will have access to other services 24 hours a day, seven days a week.
	All CRS will have access to current community contact information.
	Protocols are in place for providing referral and support (i.e., income support, dealing with employers, access to long/short-term housing, addressing family issues) based on consumer-articulated needs.

	Domain	Indicators
	Accessibility	 Access to psychiatrists and other mental health professionals Access to primary care Wait times for needed services Denial of service
	Appropriateness	 Hospital readmission rate Involuntary committal rate Length of stay in acute care Time in community programs Level of service and setting appropriate to needs of individual Availability of community services Criminal justice system involvement Community/institutional balance Treatment protocols for co-morbidity
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to the homeless Access to psychiatrists and other mental health professionals Wait times for needed services Availability of after hours care Availability of transportation Denial of service
	Competence	Resources available for on-the-job development and continuous learning
	Appropriateness	 Best practices services/supports provided to persons with SMI Level of service and setting appropriate to needs of individual Consumer/family perception of appropriateness Availability of community services Criminal justice system involvement Treatment protocols for co-morbidity
	Accessibility	 Service reach to persons with serious mental illness (SMI) Service reach to the homeless Access to psychiatrists and other mental health professionals Access to primary care Wait times for needed services Consumer/family perception of accessibility Access to continuum of mental health service Criminal justice system involvement Denial of service

Function	Standard
Review/ Follow-up/ Referral	The consumer must have the opportunity to review, discuss and comment on the service and its appropriateness.
	A review process must be established that includes an assessment of consumer outcomes including status and stability (e.g., Did the consumer perceive return of control? Was the crisis stabilized?)
	Written protocols must be established for providing referral and transition to post-crisis services. Referrals to post-crisis services must be based on consumer-articulated needs.
	CRS must ensure that staff are provided with information about other relevant community resources.
	A written follow-up plan must be developed upon completion of service, which will include criteria for follow-up, re-entry and linkage with other services.
	CRS must ensure that staff are provided with information about other relevar community resources. A written follow-up plan must be developed upon completion of service, which

	Domain	Indicators
A STEEL STATE OF THE STATE OF T	Appropriateness	Consumer/family perception of appropriateness
	Acceptability	 Consumer/family satisfaction with service received Formal complaints mechanisms in place Cultural sensitivity
	Effectiveness	 Community tenure Mortality Criminal justice system involvement Clinical status Functional status Involvement in meaningful daytime activity Housing status Quality of life Physical health status
	Appropriateness	 Best practices services/supports provided to persons with SMI Treatment protocols for co-morbidity Level of service and setting appropriate to needs of individual Consumer/family perception of appropriateness Availability of community services
	Competence	 Resources available to train staff to meet required competencies for role Resources available for on-the-job development and continuous learning
	Effectiveness	 Community tenure Criminal justice system involvement Clinical status Functional status Involvement in meaningful daytime activity Housing status Quality of life Physical health status
	Acceptability	 Consumer/family involvement in service delivery and planning Cultural sensitivity Consumer/family choice of services

Function	Standard
Monitoring and Evaluation	Consumer satisfaction (including consumers, families and outside agencies) must be monitored continuously, and the results used to make service improvements.
	A consumer satisfaction survey will be developed and implemented on an annual basis, with a target of 80% satisfaction, to the best extent possible, given the mandate of the CRS.
	All organizations and agencies must evaluate some aspect of their programs annually using best practices and published standards.

Domain	Indicators
Acceptability	 Consumer/family satisfaction with service received Consumer/family involvement in treatment decisions Formal complaints mechanisms in place Patient bill of rights Consumer/family involvement in service delivery and planning Cultural sensitivity Consumer/family choice of services
Acceptability	 Consumer/family satisfaction with service received Consumer/family involvement in treatment decisions Formal complaints mechanisms in place Patient bill of rights Consumer/family involvement in service delivery and planning Cultural sensitivity Consumer/family choice of services
Appropriateness	 Fidelity: adherence to best practices Best practices services/supports provided to persons with SMI
Effectiveness	 Community tenure Criminal justice system involvement Clinical status Functional status Involvement in meaningful daytime activity Housing status Quality of life Physical health status Mortality

Function	Standard
Information, Liaison, Advocacy, and Consultation/ Collaboration	
	A plan must be developed and published that identifies community linkages.
	The crisis worker must be knowledgeable about services that are accessible and relevant to consumers' interests in order to provide up-to-date information.
	CRS must ensure that staff is provided with information about other relevant community resources.

	Domain	Indicators
	Efficiency Continuity	 Mental health spending per capita Proportion of staff funding spent on administration and support Needs based allocation strategy Resource intensity planning tool Budget and tools for evaluation and performance monitoring Continuity mechanisms
	Effectiveness	Community follow-up after hospitalization Community tenure
		 Criminal justice system involvement Clinical status Functional status Involvement in meaningful daytime activity Housing status Quality of life Physical health status
	Competence	 Resources available to train staff to meet required competencies for role Resources available for on-the-job development and continuous learning
	Appropriateness	 Fidelity: adherence to best practices Treatment protocols for co-morbidity Level of service and setting appropriate to needs of individual Consumer/family perception of appropriateness Availability of community services Criminal justice system involvement Community/institutional balance
	Effectiveness	 Community tenure Mortality Criminal justice system involvement Clinical status Functional status Involvement in meaningful daytime activity Housing status Quality of life Physical health status

Appendix F: Literature Review – Summary of Key Research Findings on Crisis Response Services

Research literature on crisis response services (CRS) was reviewed to inform standard development. In general, there is limited research available in this area and it is under-studied compared with other areas in mental health. Of the research studies available, most were descriptive in nature and explored the approaches/components of crisis service delivery. Few studies used an experimental design to evaluate the effectiveness of CRS.

The following presents a summary of the findings of these literature reviews.

Review of Best Practices in Mental Health Reform (1997)

The 1997 Review of Best Practices in Mental Health Reform (Goering et al.) prepared for the Federal/Provincial/Territorial Advisory Network on Mental Health found that:

- Compared to other service areas such as case management, the components of CRS have been poorly studied.
- Most of the literature on CRS has been descriptive in nature, lacking an experimental design. Only one study (Merson et al., 1992) conducted a clinical study comparing inpatient hospitalization to care in the community. Other studies evaluated components of CRS including mobile crisis units (Fisher et al., 1990; Reding & Raphelson, 1995), crisis housing (Leaman, 1987; Bond et al., 1989) and hospital-based services (Gillig et al., 1989; Lambert, 1995).
- There is almost no evidence supporting the efficacy of different crisis interventions. (Geller et al., 1995)
 surveyed mobile crisis services and found that, although the services had been accepted and
 implemented, no systematic evaluations have been carried out. However, the studies reviewed suggest:
 - Crisis housing provides a viable alternative to hospitalization for persons with serious mental illness,
 - Diversion programs are effective, and
 - Crisis centres can serve persons with psychosocial problems.
- Based on these findings, the following key elements of best practice for CRS were identified:
 - Use of minimally intrusive options,
 - Programs available to divert people from inpatient hospitalization, and
 - Evaluation/research protocols are incorporated into crisis programs.

Inter-jurisdictional Review

A literature review and an inter-jurisdictional scan were conducted to assess the state of CRS standards. Literature and documents were reviewed for service standards currently in place or being designed. The review revealed that:

- There is considerable on-going work within Canada and around the world. All Canadian jurisdictions reviewed had developed or were in the process of developing services and standards (British Columbia Ministry of Health, 2003; Manitoba Department of Health, 1997; Nova Scotia Department of Health, 2003; Newfoundland Department of Health and Community Services, 2003). Australia, the United Kingdom and the United States were at various points in developing standardized programs and services. Australia and the United Kingdom do not have standards specific to CRS but have developed national standards that are applied to all mental health services (Commonwealth of Australia, 1996; National Health Service, 1999).
- There is a lack of firm standards that are evidence-based, or that have been evaluated.
- Existing literature is limited and tends to evaluate services rather than evaluate the effectiveness
 of crisis interventions, although Cochrane et al. (1997) found that CRS are effective in diverting
 people from inpatient hospitalization and for minimizing the intrusiveness of the intervention.

- There is consensus that CRS are an essential component of the overall mental health system;
 however, it was found that definitions and implementation of services varied across jurisdictions.
- The Community Mental Health Evaluation Initiative conducted a series of evaluations involving various types of mental health programs, including a one-year evaluation of two mobile crisis programs.
 The findings from the evaluation of the two mobile crisis programs suggest the need for:
 - Integrating mobile crisis services with other mental health services
 - Clear lines of accountability and responsibility and attention to quality assurance
 - Close supervision and feedback to mobile crisis workers
 - A clear definition of the target population
- Educational and communication opportunities among mobile crisis services and other agencies (i.e., policy, psychiatric services within a coordinated network of mental health services). (Ferris et al., 2000)

Literature Review: Standards Validation

A literature review was conducted to validate the service standards for CRS. Levels of Evidence were used to review the literature supporting the standards.

Levels of Evidence

- A four-level typology of evidence adapted from the typology developed by the Nova Scotia
 Department of Health (Nova Scotia Department of Health, 2003) was used to assess the literature
 against the proposed crisis response service standards.
- Level 1 involved direct evidence of effectiveness (i.e., specific standard evaluated as independent variable in a study and shown to produce positive outcomes).
- Level 2 considered indirect evidence of effectiveness (i.e., specific standard is one of the characteristics or ingredients of a program which has been shown to be effective).
- Level 3 involved studies based on expert opinion/consensus of effectiveness or correlational evidence of standards being associated with positive consumer outcomes.
- Level 4 involved expert opinion (i.e., defined by the working group or other Canadian jurisdictions in absence of empirical support in research literature).

Review Findings

Overall, there were limited experimental research studies supporting the development of standards. In general, the research tended to be descriptive but did not identify specific program components or best practices. Evidence for the proposed standards was largely based on expert stakeholder opinion. Other Canadian jurisdictions were found to have developed similar standards based on similar levels of evidence (i.e., Nova Scotia, British Columbia).

Standard	Evidence for Standard
 Upon identification of a crisis, the first contact with the consumer by the crisis response service (CRS) must be established within 90 minutes. A crisis requiring in-person contact will be responded to as soon as 	 Level 4 – Based on expert opinion. British Columbia (BC) – Standard for mobile crisis units requires immediate response to crisis. Level 4 – Based on expert opinion.
possible. Response time should be within 24 hours, with consideration for travel time, weather, etc.	BC – Standard for mobile crisis units requires the development of guidelines for the prompt determination of a response strategy to crisis calls.
 Crisis support telephone lines must be configured to include a queuing system that lasts no longer than 15 minutes. During the wait time, voice instructions for alternative crisis management options must be provided (e.g., 911, local emergency number, another crisis line). 	• Level 4 – Based on expert opinion.
 Protocols must be in place with related service providers (e.g., case management services, psychiatrists, hospitals, primary care teams, etc.) in order to ensure access to necessary medical, psychiatric and psychological/social assessments and existing crisis management protocols. If access to other health records will assist in planning with the consumer, documented consent is required. 	 Level 2 – Three studies. Found participation of psychiatrist (Reding & Raphelson, 1995) and the ability to prescribe medication (Tufnell et al., 1985; Reynolds et al., 1990) critical to mobile crisis teams in reducing hospitalization. Level 4 – Based on expert opinion. Nova Scotia (NS) – Standard for the availability of timely consultation to various service providers to assist in identifying and intervening in emergencies.
	Manitoba – Identifies multidisciplinary service delivery as a key value of the mental health system as a whole (including crisis response services). Newfoundland – Identifies the provision of CRS through partnership of primary care health teams and mobile crisis
	first contact with the consumer by the crisis response service (CRS) must be established within 90 minutes. • A crisis requiring in-person contact will be responded to as soon as possible. Response time should be within 24 hours, with consideration for travel time, weather, etc. • Crisis support telephone lines must be configured to include a queuing system that lasts no longer than 15 minutes. During the wait time, voice instructions for alternative crisis management options must be provided (e.g., 911, local emergency number, another crisis line). • Protocols must be in place with related service providers (e.g., case management services, psychiatrists, hospitals, primary care teams, etc.) in order to ensure access to necessary medical, psychiatric and psychological/social assessments and existing crisis management protocols. If access to other health records will assist in planning with the consumer, documented consent

Function	Standard	Evidence for Standard
Assessment and Planning (continued)		BC – Standards were developed based on the principle that CRS require members to function as interdisciplinary teams and develop and maintain service links with other CRS providers. Defines assessments that identify and integrate range of factors (biological, psychological and/or social) as a key principle of CRS. Also has a standard for its mobile crisis units and walk-in crisis stabilization services that require guidelines to assess risk, presence of mental illness, stressors and the need for medical evaluation.
	Services must be provided in the consumer's place of choice wherever possible.	Level 4 – Based on expert opinion. Manitoba – Identifies this as a key value of the mental health system as a whole (including crisis response services).
	The CRS will ensure that staff have training and core competencies in a variety of areas, to the best extent possible, (e.g., risk assessment; suicide assessment; crisis intervention; safety standards; conflict resolution; anti-racism training; psychiatric symptomatology and psychiatric medications).	Level 4 – Based on expert opinion. NS – Requires CRS providers to have core competencies in risk assessment. Manitoba – Recognizes training as a key policy direction to strengthen the mental health system. BC – Requires staff to be trained to assess risk, mental illness, stressors and the need for medical evaluation. Also identifies CRS staff skills including: knowledge of psychiatric illness and range of psychiatric conditions, suicide assessment and management, assaultive behaviour and acute psychosis, crisis intervention skills, comprehensive mental status exam, etc. Risk management is identified as a key principle of CRS.
	 The CRS must ensure that workers are provided with information regarding relevant legislative reporting requirements. 	Level 4 – Based on expert opinion. BC – Requires mobile and hospital based crisis teams to have training in the Mental Health Act and relevant legislation.

Function *	Standard	Evidence for Standard	
Crisis Support/ Counseling	Services are consumer directed and will be provided in the least intrusive manner possible.	 Level 4 – Based on expert opinion. Manitoba – Recognizes consumer centredness as a key policy direction to strengthen the mental health system. Newfoundland – Defines the involvement of the individual and caregiver in planning and decision making as a Strategic Direction. 	
	Written protocols for consumer and service provider safety must be established.	 Level 4 – Based on expert opinion. BC – Defines risk management as a key principle of CRS and sets standards to address the safety of consumers and staff. 	
	 Written protocols must be developed to ensure that timely consultation is available with various service providers (e.g., physicians, guidance counsellors, CAS workers) to assist in identifying and intervening in actual emergencies; documented consent is required. 	 Level 4 – Based on expert opinion. NS – Identifies this as standard for its CRS. Manitoba – Identifies multidisciplinary service delivery as a key value of the mental health system as a whole (includes CRS). 	
	Short-term crisis support/counseling will be available to provide risk assessment, de-escalation and safety planning, to the best extent possible, given the mandate of the CRS.	Level 4 – Based on expert opinion. BC – Requires walk-in crisis services to provide crisis intervention to help consumers make sense of the crisis, mobilize personal coping resources and formulate a plan to cope more effectively.	

Function	Standard.	Evidence for Standard
Medical Intervention	The CRS must have written protocols for initiating and accessing medical interventions in a timely manner.	 Level 4 – Based on expert opinion. NS – Set the standard that a policy will identify situations and circumstances where medical clearance/assessment is required. BC – Has a standard requiring consultation regarding medical/medication issues.
	The CRS will have procedures/ guidelines for responding to emergency and non-emergency medical situations.	 Level 4 – Based on expert opinion. BC – Requires staff to consult regarding medical/medication issues and receive training in assessing for the need for medical evaluation.
Environmental Intervention and Crisis Stabilization	All CRS will have access to other services 24 hours a day, seven days a week.	 Level 4 – Based on expert opinion. Saskatchewan, Quebec and Nova Scotia have set the standard of 24/7 access to crisis services. Newfoundland requires the 24-hour availability of crisis services provided through its primary care team. All view CRS as part of a continuum of mental health services.
	 All CRS will have access to current community contact information. Protocols are in place for providing referral and support (i.e., income support, dealing with employers, access to long/short-term housing, addressing family issues) based on consumer-articulated needs. 	 Level 4 – Based on expert opinion. Level 4 – Based on expert opinion. BC – Identified access to a range of follow-up services as a key principle of CRS. Set standard that crisis team is engaged until links with appropriate services are made and that a clear, well-documented care plan and discharge plan must be developed.

Function	Standard	Evidence for Standard
Review/ Follow-up/ Referral	The consumer must have the opportunity to review, discuss and comment on the service and its appropriateness.	Level 4 – Based on expert opinion. Newfoundland, Nova Scotia and Manitoba all identified the inclusion of the consumer in evaluating services.
	A review process must be established that includes an assessment of consumer outcomes including status and stability (e.g., Did the consumer perceive return of control? Was the crisis stabilized?)	Level 4 – Based on expert opinion. BC – Standard for its Community Crisis Stabilization Services that discharge planning will include family and key supports. NS – Generic standards for all mental
		health services includes mutually determining goals/outcomes and revising and evaluating the treatment plan and services.
	Written protocols must be established for providing referral and transition to post-crisis services. Referrals to post-crisis services must be based on consumer-articulated needs.	Level 4 – Based on expert opinion. BC – Identified access to a range of follow-up services as a key principle of CRS. Set standard that crisis team is engaged until links with appropriate services are made and a clear, well-documented care plan and discharge plan is developed.
		NS – Generic standards for all mental health services includes developing treatment and discharge plans that include appropriate linkage and coordination with other services and resources.
	CRS must ensure that staff are provided with information about other relevant community resources.	Level 4 – Based on expert opinion. NS – Standard for providing continuity of services through coordination with other community crisis services. Manitoba – Defines a key value of the
		mental health system as working in partnership with consumers, service providers and government in the planning and delivery of service.

Function	Standard	Evidence for Standard
Review/ Follow-up/ Referral (continued)		Newfoundland – One of its strategic directions is to ensure there are connections to the community services people need.
	A written follow-up plan must be developed upon completion of service, which will include criteria for follow-up, re-entry and linkage with other services.	 Level 4 – Based on expert opinion. BC – Access to a range of follow-up services is a key principle of CRS. Set standard that crisis team is engaged until links with appropriate services are made and that a clear, well-documented discharge plan must be developed.
		NS – Generic standards for all mental health services include development and monitoring of individualized follow-up plan.
Monitoring and Evaluation	Consumer satisfaction (including consumers, families and outside agencies) must be monitored continuously, and the results used to make service improvements.	 Level 4 – Based on expert opinion. Newfoundland – Key strategic direction for entire mental health system considers accountability and requires the inclusion of consumers and families in service evaluation.
	 A consumer satisfaction survey will be developed and implemented on an annual basis, with a target of 80% satisfaction, to the best extent possible, given the mandate of the CRS. 	 Level 4 – Based on expert opinion. Newfoundland – Key strategic direction for entire mental health system considers accountability and requires the inclusion of consumers and families in service evaluation.
	All organizations and agencies must evaluate some aspect of their programs annually using best practices and published standards.	 Level 4 – Based on expert opinion. NS – Generic standards for all mental health services includes annual monitoring of compliance with standards and utilization of services. Newfoundland – Key strategic direction for entire mental health system considers accountability and requires the inclusion of consumers and families in service evaluation.

Function	Standard	Evidence for Standard.
Information, Liaison, Advocacy, and Consultation/ Collaboration	Service providers must identify gaps in service and develop means for collaborating with other relevant community resources in order to meet unmet needs.	 Level 4 – Based on expert opinion. NS – Standard for providing continuity of services through coordination with other crisis services in the community. Manitoba – Defines a key value of the mental health system as working in partnership with consumers, service providers and government in the planning and delivery of service. Newfoundland – Identifies a strategic direction of establishing a continuum of mental health services (including crisis services) linked with primary care teams.
	A plan must be developed and published that identifies community linkages.	 Level 4 – Based on expert opinion. NS – Generic standards for all mental health services includes the development of a treatment and discharge plan that incorporates other services and resources. BC – Identified access to a range of follow-up services as a key principle of CRS. Set standard that crisis team is engaged until links with appropriate services are made and that a clear, well-documented treatment and discharge plan is made.
	The crisis worker must be knowledgeable about services that are accessible and relevant to consumers' interests in order to provide up-to-date information.	 Level 4 – Based on expert opinion. Manitoba and Newfoundland identified the inclusion of family and their access to supports as key. BC – Requires the communication of pertinent information to consumers and family members.
	 CRS must ensure that staff is provided with information about other relevant community resources. 	• Level 4 – Based on expert opinion.

Appendix G: Bibliography

Bond, G.R., Witheridege, T.E., Wasmer, D., Dincin, J., McRae, S.A., Mayes, J. & Ward, R.S. (1989). A comparison of two crisis housing alternatives to psychiatric hospitalization. *Hospital and Community Psychiatry*, 40(2), 177-183.

British Columbia Ministry of Health (2003). B.C.'s Mental Health Reform Best Practices: Crisis Response/Emergency Services. Retrieved on the World Wide Web on July 23, 2004 at http://www.healthservices.gov.bc.ca/mhd/pdf/bp_crisis_response.pdf

California Mental Health Planning Council (2003). California Mental Health Master Plan: A Vision for California. California Department of Mental Health. Retrieved on the World Wide Web on October 18, 2004 at http://www.dmh.ca.gov/mhpc/docs/Master%20Plan/mstrplan.pdf

Cochrane, J., Durbin, J., & Goering, P. (1997). Best Practices in Mental Health Reform – Discussion Paper. Prepared for the Federal/Provincial/Territorial Advisory Network on Mental Health.

Commission on the Future of Health Care in Canada (2002). Final Report: Building on Values: The Future of Health Care in Canada. Retrieved on the World Wide Web on Sept. 9, 2004 at http://www.hc-sc.gc.ca/english/care/romanow/hcc0086.html

Commonwealth of Australia. (1996). National Standards for Mental Health Services. Retrieved October 20, 2004 on the World Wide Web at http://www.health.gov.au/internet/wcms/Publishing.nsf/Content/mentalhealth-mhinfo-standards-nsmhs.htm

Ferris, L.E., Shulman, K., Williams, J., Desiato, C., Sandercock, J., & Mills, W. (2000). A Descriptive Evaluation of Three Ontario Mobile Crisis Programs for Clients with Severe and Persistent Mental Illness (SPMI). Unpublished report for the Community Mental Health Evaluation Initiative.

Fisher, W.H., Geller, J.L., & Wirth-Cauchon, J. (1990). Empirically assessing the impact of mobile crisis capacity on state hospital admission. *Community Mental Health Journal*, 26(3), 245-253.

Geller, J.L., Fisher, W.H., & McDermeit, M. (1995). A national survey of Mobile Crisis Services and their evaluation. *Psychiatric Services*, 46(9), 893-897.

Gillig, P.M., Hillard, J.R., Bell, J., Combs, H.E., Martin, C. & Deddens, J.A. (1989). The psychiatric emergency service holding area: Effect on utilization of inpatient resources. *American Journal of Psychiatry*, 146(3), 369-372.

Goering, P., Boydell, K., Butteril, D., Cochrane, J., Durbin, J., Rogers, J. & Trainor, J. (1997). *Review of Best Practices in Mental Health Reform.* Prepared for the Federal/Provincial/Territorial Advisory Network on Mental Health.

Health Canada (2003). 2003 First Ministers' Accord on Health Care Renewal. Retrieved on the World Wide Web on September 9, 2004 at http://www.hc-sc.gc.ca/english/hca2003/accord.html

Lambert, M. (1995). Psychiatric crisis intervention in the general emergency service of a Veterans Affairs hospital. *Psychiatric Services*, 46(3), 283-284.

Leaman, K. (1987). A hospital alternative for patients in crisis. *Hospital and Community Psychiatry*, 38(11), 1221-1223.

Manitoba Department of Health (1997). Core Health Services in Manitoba. Manitoba Health. Retrieved on World Wide Web on July 22, 2004 at http://www.gov.mb.ca/health/rha/core.pdf

Merson, S., Tyrer, P., Onyett, S., Lack, S., Brikett, P., Lynch, S. & Johnson, T. (1992). Early intervention in psychiatric emergencies: a controlled clinical trial. *The Lancet*, 339, 1311-1314.

National Health Service (1999). National Service Framework for Mental Health: Modern Standards and Service Models. United Kingdom.

Newfoundland Department of Health and Community Services. (2003). Working together for Mental Health: a proposed mental health services strategy for Newfoundland and Labrador. Retrieved on World Wide Web on July 23, 2004 at http://www.gov.nf.ca/health/publications/pdfiles/Mental%20Health%20Strategy%20Disc%20Doc%20Nov%202003.pdf

Nova Scotia Department of Health. (2003). Standards for Mental Health Services in Nova Scotia. Retrieved on World Wide Web on July 23, 2004 at www.gov.ns.ca/health/downloads/Standards.pdf

Ohio Department of Mental Health, Office of Program Evaluation and Research (1999). Towards Best Practices: Top Ten Findings from the Longitudinal Consumer Outcomes Study 1999. Columbus, Ohio, Ohio Department of Mental Health, Office of Program Evaluation and Research.

Oklahoma Department of Mental Health and Substance Abuse Services. (2003). Chapter 23: Standards and Criteria for Community-Based Structured Crisis Centers. Retrieved on World Wide Web on July 24, 2004 at http://www.odmhsas.org/eda%20II/Title450-23.pdf

Ontario Government (2004). 2004 Ontario Budget Backgrounder Transforming Health Care. Retrieved on the World Wide Web on September 9, 2004 at http://www.gov.on.ca/FIN/bud04e/bke1.htm

Ontario Ministry of Health and Long-Term Care (2004). *Public Information – Taking Action to Improve Health Care*. Retrieved on the World Wide Web October 18, 2004 at http://www.health.gov.on.ca/transformation/#1

Ontario Ministry of Health and Long-Term Care (2003). Mental Health Accountability Framework.

Ontario Ministry of Health and Long-Term Care (1999). Making It Happen: Implementation Plan for Mental Health Reform.

Ontario Ministry of Health and Long-Term Care (1999). Making It Happen: Operational Framework for the Delivery of Mental Health Services and Supports.

Reding, G.R., & Raphelson, M. (1995). Around-the-clock mobile psychiatric crisis intervention. *Community Mental Health Journal*, 31(2), 179-190.

Reynolds, I., Jones, J.E., Berry, D.W., et al. (1990). A crisis team for the mentally ill: the effect on patients, relatives, and admissions. *Medical Journal of Australia*, 152, 646-652.

State of Vermont Department of Developmental and Mental Health Services (2002). *The Statewide System of Care Plan for Adult Mental Health in Vermont: Fiscal Years 2002-2004*. Retrieved on the World Wide Web on July 22, 2004 at http://www.state.vt.us/dmh/docs/publications/system-of-care-plans.html#adult

Tufnell, G., Bouras, N., Watson, J.P., et al. (1985). Home assessment and treatment in a community psychiatric service. *Acta Psychiatrica Scandinavica*, 72, 20-28.



♥ Ontario

FNIHB Manitoba Region - Mental Wellness Team - Psychosocial Response to Community Crisis or Disaster

